

Evaluation of endophytic aquatic hyphomycetes for their antagonistic activity against pathogenic bacteria

Pratibha Arya^{1*} and S. C. Sati²

¹Botany Department, Govt. P. G. College Augustyamuni, Rudraprayag, Garhwal, Uttarakhand.

²Botany Department, Kumaun University, Nainital, Uttarakhand.

Evaluation of some riparian endophytic aquatic hyphomycetous fungi viz., *Heliscus lugdunensis*, *Tetrachaetum elegans*, *Tetracladium marchalianum*, *T. breve* and *T. nainitalense* from surface sterilized roots of healthy riparian plants has been carried out for their possible antibacterial activity. Pathogenic test bacteria viz., *Agrobacterium tumefaciens*, *Bacillus subtilis*, *Erwinia chrysanthemi*, *Escherichia coli* and *Xanthomonas phaseoli* were used. The antibacterial test was conducted by using agar-disc technique. Two aquatic hyphomycetes, *T. elegans* and *T. marchalianum* showed significant antagonistic activity against test bacteria.

Keywords: Antibacterial activity, root endophytes, *Tetrachaetum elegans*, *Tetracladium marchalianum*.

INTRODUCTION

Endophytes are microorganisms that inhabit the plant tissues in their life cycle without causing any apparent harm to their host (Petrini, 1991). Their presence implies a symbiotic interaction with the host plants (Azevedo *et al.* 2003). It has been known that endophytic fungi are important sources of bioactive compounds (Strobel, 2003; Pan *et al.*, 2008). A number of new bioactive compounds from endophytes have been recognized as potential sources of antimicrobial substances (Strobel, 2003; Li *et al.*, 2005).

Uses of microorganisms or their metabolites to prevent diseases offer an attractive alternative or supplement to disease management without the negative impact of chemical control (Gani and Ganesh, 2009). Many of the aquatic hyphomycetes have now been reported as root endophytes (Fisher *et al.*, 1991; Sridhar *et al.*, 1992; Sridhar and Raviraja, 1995; Sati and Belwal, 2005; Arya and Sati, 2010). Occurrence of aquatic hyphomycetes as root endophytes in healthy plants indicates that they may have beneficial role in plant health (Bills and Polishook, 1992; Dreyfus and Chapela, 1992; Singh and Waingankar, 2005). Endophyte is also well known to constitute a valuable source of secondary metabolites for the discovery of new potential therapeutic drugs (Miller,

1995). Recently, Sati and Arya (2010 a) reported the significant effects of two aquatic hyphomycetes viz., *Heliscus lugdunensis* and *Tetrachaetum elegans* in plant growth in pot experiments.

Interaction of aquatic hyphomycetes with bacteria and terrestrial fungi has been documented previously (Chamier *et al.*, 1984; Gulis and Suberkropp, 2003). Platas *et al.* (1998) and Gulis and Stephanovich (1999) demonstrated the antagonistic activity of aquatic hyphomycetes due to release of diffusible inhibitory substances. According to Gloer (1995) the secondary metabolites of aquatic hyphomycetes could result in the discovery of new natural bioactive products of medicinal and agricultural importance. 'Quinapathin' from the aero-aquatic hyphomycetes (*Helicoon richonis* (Boud.) Linder (Fisher *et al.*, 1988; Adriaenssens *et al.*, 1994) and 'Anguillosporal', from *Anguillospora longissima* has resulted in the discovery of new metabolite (Harrigan *et al.*, 1995). The earlier studies on antagonistic activity of ectomycorrhizal fungi and other endophytic fungi have shown positive results (Raviraja *et al.*, 2006; Gbolagade *et al.*, 2007; Vaidya *et al.*, 2005; Gulis and Stephanovich, 1999).

Intra and interspecific interaction of aquatic hyphomycetes in relation to aquatic ascomycetes and release of diffusible inhibitory substances has also been reported (Shearer and Zare-Maivan, 1988; Barlocher, 1991). Sati and Arya (2010 b) also reported the

antagonistic effects of some root endophytic aquatic hyphomycetes against different plant pathogenic fungi.

The purpose of this paper was to study the antagonistic activity of five Aquatic Hyphomycetes viz., *Heliscus lugdunensis*, *Tetrachaetum elegans*, *Tetracladium breve*, *T. marchalianum* and *T. nainitalense* isolated from the roots of riparian plants against pathogenic bacteria (plants as well as animals).

MATERIALS AND METHODS

Isolation of riparian root endophytic aquatic hyphomycetes:

Roots of healthy riparian plants were collected from ravine areas near Nainital, Kumaun Himalaya, India (29.39° N 79.45° E). Root pieces were harvested, washed in sterile water, and surface sterilized by immersing it into 0.01% sodium hypochlorite solution (3-6 min) and then in 96% ethanol (30 s) following Fisher et al. (1991). Root pieces (1-2 cm) were then placed in 2% malt extract agar Petri dishes (90 mm diameter) and incubated at 20±2°C for 10-15 days in dark. Malt extract plates were also supplemented with 0.5 g/L Streptomycin to suppress the bacterial contamination. Isolation of Aquatic Hyphomycetous fungi was done from the hyphae that grew in agar (Figure 1). The isolates were identified with the help of relevant monographs and papers (Ingold, 1975; Marvanova, 1997). *Heliscus lugdunensis* Sacc. and Therry was isolated from *Strobilanthes alatus*, *Tetrachaetum elegans* Ingold from *Pilea scripta*, *Tetracladium breve* Roldan from *Eupatorium adenophorum*, *T. marchalianum* De Wildeman from *Geranium nepalense* and *T. nainitalense* Sati & Arya from *Eupatorium adenophorum*.

Antagonistic activity against pathogenic bacteria

Antagonistic activity of aquatic hyphomycetes against pathogenic test bacteria viz., *Erwinia chrysanthemi*, *Xanthomonas phaseoli*, *Agrobacterium tumefaciens*, *Bacillus subtilis* and *Escherichia coli* was tested by using agar-disc technique. Bacterial species were grown in Nutrient Agar broth (beef extract = 1 g, yeast extract (oxid) = 2 g, peptone (bacteriological) = 5 g, sodium chloride = 5 g and distilled water = 1 l). Nutrient agar broth was seeded with test bacteria at 37 ± 2°C for 48 h to prepare the bacterial suspension.

The mycelial disks (7 mm diam.) of actively growing colonies of aquatic hyphomycetes were cut from the periphery of the culture plates and aseptically placed in the centre of the assay plates (Figures 2-3). These assay plates were prepared by 15 ml Nutrient Agar (NA) medium poured in 90 mm Petri plates. Nutrient Agar surface was seeded with bacterial suspension of A.




Figure 1. Culture plates of different Aquatic Hyphomycetes. a: *Heliscus lugdunensis*; b: *Tetrachaetum elegans*; c: *Tetracladium breve*; d: *T. marchalianum*; e: *T. nainitalense*.

tumefaciens, *E. coli*, *B. subtilis*, *E. chrysanthemi* and *X. phaseoli*. This experiment was performed in three replicates and the plates with bacterial growth without the aquatic hyphomycetes were used as control. Plates were inspected after 24 h of incubation for the presence of clear inhibitory zones around the agar disks indicating the antagonistic activity of the aquatic hyphomycetes used in this experiment.

Analysis of data

One way Analysis of Variance (ANOVA) was performed for *T. elegans* and *T. marchalianum*. *T. elegans* showed significant variations ($P < 0.001$) while *T. marchalianum* was found not significant in antagonistic activity against bacteria.

Table 1. Inhibition of bacterial growth (mm) by Aquatic Hyphomycetes (\pm SEM based on 3 replicates).

Fungal Isolates	Inhibition (mm) of Test Bacteria				
	<i>A. tumefaciens</i> (Gram-ve)	<i>E. coli</i> (Gram-ve)	<i>X. phaseoli</i> (Gram-ve)	<i>B. subtilis</i> (Gram +ve)	<i>E. chrysanthemi</i> (Gram -ve)
<i>T. elegans</i>	9.0 \pm 0.00	8.33 \pm 0.33	8.0 \pm 1.15	8.67 \pm 0.67	—
<i>H. lugdunensis</i>	—	—	—	—	—
<i>T. marchalianum</i>	8.0 \pm 0.00	12 \pm 2.00	9.33 \pm 0.33	8.33 \pm 0.88	8.67 \pm 0.33
<i>T. breve</i>	—	—	—	—	—
<i>T. nainitalense</i>	—	—	—	—	—

— = No activity.




Figure 2 a-d. Inhibition of plant pathogenic bacteria by *Tetrachaetum elegans*: a-*A. tumefaciens*; b-*E.coli*; c-*X.phaseoli*; d-*B.subtilis*.




Figure 3 a-e. Inhibition of plant pathogenic bacteria by *Tetracladium marchalianum*: a-*A. tumefaciens*; b-*E.coli*; c-*X.phaseoli*; d-*B.subtilis*; e-*E. chrysanthemi*.

RESULTS

Activity against pathogenic bacteria

The result of antagonistic activity against five pathogenic bacteria is presented in Table 1. It is evident that out of the five fungal species taken in the current study, *T. elegans* showed activity against four pathogenic bacteria namely *A. tumefaciens*, *E. coli*, *X. phaseoli* and *B. subtilis* (Figure 2, a-d) showing a clear zone of inhibition (Table 1). *T. marchalianum* showed its activity against all the test bacteria viz., *A. tumefaciens*, *E. coli*, *X. phaseoli*, *B.*

subtilis and *E. chrysanthemi* (Figure 3, a-e). However, *H. lugdunensis*, *T. breve* and *T. nainitalense* showed no inhibitory effect against all the test bacteria (Table 1).

DISCUSSION

In this study the antagonistic activity of aquatic hyphomycetes against pathogenic bacteria was screened by using agar disc technique. In a test performed by Gulis and Stephanovich (1999), they observed *H. lugdunensis* as biologically inactive, showing no activity against Gram

-ve and Gram +ve bacteria, yeast and Hyphomycetes in "agar well" technique using its metabolite. It is interesting to note that in the present screening test also *H. lugdunensis* showed no antagonistic activity against pathogenic bacteria (Table 1). In the present experiment, *T. marchalianum* showed activity against all the test bacteria (Table 1). This supports the work of Gulis and Stephanovich (1999).

Gulis and Suberkropp (2003) screened 28 isolates of aquatic hyphomycetes belonging to *Alatospora acuminata* (6 isolates), *Anguillospora filiformis* (5 isolates), *Articulospora tetracladia* (10 isolates), *Tetrachaetum elegans* (4 isolates) and *Tricladium chaetocladium* (3 isolates) against 16 bacterial isolates. These aquatic hyphomycetes inhibited the bacterial growth by forming the clear zones in the bacterial loans around wells containing the fungal culture broth. In the present study *T. elegans* also showed inhibitory activity against 4 test bacteria (Figure 2, Table 1). It is worth mentioning here that all the previous studies on antibacterial activity were determined by using metabolites of endophytic aquatic hyphomycetes but the present investigation was conducted to determine the antagonistic role of these fungi by using agar disc technique for the first time.

CONCLUSION

Aquatic hyphomycetes occurring as endophytes not only help in developmental and physiological activity of plants but also antagonize their bacterial pathogens. Endophytic aquatic hyphomycetes with potential as antibacterial can be used in pharmaceutical companies for the large scale production of useful compounds.

REFERENCES

- Adriaenssens P, Anson E, Begley MJ, Fisher PJ, Orrel KG, Webster J, Whitehurst JS (1994). Quinaphthin, a metabolite produced by *Helicoon richonis*. *J. Chem. Society. Perkin Transactions 1*(14): 2007-2010.
- Arya P, Sati SC (2010). Four species of Aquatic Hyphomycetes occurring as New Root Endophytes. *Natl. Acad. Sci. Lett.* 33: 299-301.
- Azevedo JL, Maccheroni WJr, Pereira PO, Arajio JL (2003). Endophytic microorganism: A review on insect control and recent advances on tropical plants. *EJB: Electronic J. Biotechnol.*
- Barlocher F (1991). Intraspecific Hyphal Interactions among Aquatic Hyphomycetes. *Mycol.* 83: 82-88.
- Bills GF, Polishook JD (1992). Recovery of endophytic fungi from *Chamaecykaris thyoides*. *Sydowia*, 44: 1-12.
- Chamier AC, Dixon PA, Archer SA (1984). The spatial distribution of fungi on decomposing alder leaves in a freshwater stream. *Oecol.* 64: 92-103.
- Daybas RA (1984). Avemectins their chemistry and pesticidal activities. En : Miyamoto J, Keaney PC (Eds.) *Pesticide chemistry* New York, Pergamon Press, 83-90.
- Dreyfus MM, Chapela IH (1992). The potential of fungi in discovery of novel, low molecular weight Pharmaceuticals. In the Discovery of Novel Natural Products with Therapeutic Potential, Butterworth Publications, Biotechnology Series, London.
- Fisher PJ, Anson AE, Webster J (1988). Quinaphthin, a new antibiotic, produced by *Helicoon richonis*. *Trans. Brit. Mycol. Soc.*, 90: 499-502.
- Fisher PJ, Petrini O, Webster J (1991). Aquatic hyphomycetes and other fungi living aquatic and terrestrial roots of *Alnus glutinosa*. *Mycol. Res.*, 95: 543- Sridhar, K.R., Chandrasekhar, K.R. and Kaveriappa, K.M. 1992. Research on the Indian Subcontinent. In "The Ecology of Aquatic Hyphomycetes (ed. Bärlocher, F.) Springer-Verlag, Heidelberg. pp.182-211.
- Gani SB, Ganesh K (2009). Preliminary screening of endophytic fungi from medicinal plants in India for antimicrobial and antitumour activity. *International journal of Pharmaceutical sciences and Nanotechnology*, 2: 566-571.
- Gbolagade J, Kigigha L, Ohimain E (2007). Antagonistic effects of extracts of some Nigerian higher fungi against selected pathogenic microorganisms. *Ame.-Eur. J. Agric. and Environ. Sci.*, 4: 364-368.
- Gloer JB (1995). Bioactive metabolite from aquatic fungi. In The VI International Marine Mycological Symposium (Incorporating Society, 8-15, July 1995, Programme) Abstracts, University of Portsmouth: Engl. p. 65
- Gulis V, Stephanovich AI (1999). Antibiotic effects of some aquatic hyphomycetes. *Mycol. Res.*, 103: 111-115.
- Gulis V, Suberkropp K (2003). Effect of inorganic nutrients on relative contributions of fungi and bacteria to carbon flow from submerged decomposing leaf litter. *Microbial Ecol.* 45: 11-19
- Harrigan GG, Armentrout BL, Gloer JB, Shearer CA (1995). New bioactive natural products from two *Anguillopsora* species. In The VI International Marine Symposium (Incorporating Freshwater Mycology). A Meeting of the British Mycological Society, 8-15 July, 1995. Programme and Abstracts, University of Portsmouth., England pp. 135.
- Ingold CT (1975). An illustrated guide to aquatic and water borne hyphomycetes (Fungi Imperfecti) with notes on their biology. *England Freshwater Biol. Assoc. Scient. Publ.* 30:96.
- Li Y, Song YC, Liu JY, Ma YM, Tan RX (2005). Anti-Helicobacter pylori substances from endophytic fungal cultures. *World J. Microbiol. Biotechnol.* 21: 553-558.
- Marvanova L (1997). Freshwater hyphomycetes: A survey with remarks on tropical taxa *In Tropical Mycology* (eds. K. K. Janardhanan, C. Rajendran, K. Natrajan and D. L. Hawksworth) Science Publishers Inc., pp.169-226.
- Miller SL (1995). Functional diversity in fungi. *Can. J. Bot.*, 73 (Suppl.1), S50-S57.
- Pan JH, Jones EBG, She ZG, Pang JY, Lin YC (2008). Review of bioactive compounds from fungi in the South China Sea. *Bot. Mar.* 51: 179-190.
- Petrini (1991). Fungal endophytes of tree leaves. In : *Microbial ecology of leaves* (edn) by J. H. Andrews and S. S., Hiran, Springer Verlag. New York, USA.
- Platas G, Pelaez F, Collado J, Villuendas G, Diez MT (1998). Screening of antimicrobial activities by aquatic hyphomycetes cultivated on various nutrient sources. *Cryptogam Mycol.* 19: 33-43.
- Raviraja NS, Maria GL, Sridhar KR (2006). Antimicrobial evaluation of endophytic fungi inhabiting medicinal plants of the Western Ghats of India. *Eng. Life Sci.*, 6: 515-520.
- Sati SC, Arya P (2010 a). Assessment of root endophytic aquatic Hyphomycetous fungi on plant growth. *Symbiosis.* 50: 143-149.
- Sati SC, Arya P (2010 b). Antagonism of some aquatic Hyphomycetes against plant pathogenic fungi. *The Scientific World J.* 10: 760-765.
- Sati SC, Belwal M (2005). Aquatic hyphomycetes as endophyte of riparian plant roots. *Mycol.* 97: 45-49.
- Shearer CA, Zare-Maivan H (1988). In vitro hyphal interactions among wood and leaf-inhabiting Ascomycetes and Fungi Imperfecti from freshwater habitats. *Mycol.* 80: 31-37.
- Singh SK, Waingankar V (2005). Endophytic fungi from medicinal plant *Holarrhena antidysenterica* and their potential antibacterial activity in Microbial diversity (Opportunities and Challenges) (Eds. S. P. Gutum et al.) from pp. 168-178.
- Sridhar KR and Raviraja NS (1995). Endophytes- a crucial issue. *Curr. Sci.*, 69: 570-571.

Strobel GA (2003). Endophytes as sources of bioactive products. *Microbe Infect.* 5:535-544.

Turhan G, Gossmann F (1994). Antagonistic activity of five *Myrothecium* species against fungi and bacteria in vitro. *J. Phytopathol.* 140: 97-113.

Vaidya GS, Shrestha K, Wallander H (2005). Antagonistic study of ectomycorrhizal fungi isolated from Baluwa Forest (Central Nepal) against with pathogenic fungi and bacteria. *Sci. World*, 3: 49-52.